

Lumiprobe Corporation

201 International Circle, Suite 135 Hunt Valley, Maryland 21030 USA

Phone: +1 888 973 6353

Fax: +1 888 973 6354 Email: order@lumiprobe.com

Biotin-11-UTP

http://www.lumiprobe.com/p/biotin-11-utp

Biotin-11-UTP is a substrate for RNA-polymerases SP6, T3, and T7. Biotinylated triphosphate is an analog of natural UTP and can be used for RNA labeling in transcription *in vitro*.

Biotinylated RNA can be used instead of radioactive-labeled RNA in many applications, including Northern and Southern blots, hybridization *in situ*, and microarray analysis. Biotinylated RNA is detected using various methods with streptavidin conjugates, and the long linker of 11 atoms allows efficient interaction of biotin with streptavidin.

Structure of Biotin-11-UTP

General properties

Appearance: colorless solid Molecular 894.48 weight:

Molecular C₂₈H₄₃N₆Li₃O₁₈P₃S formula:

 $IUPAC \\ ((2R,35,4R,5R)-5-(2,4-dioxo-5-(3-(6-(5-((3a5,45,6aR)-2-oxohexahydro-1H-thieno[3,4-d])midazol-4-yl)pentanamido) \\ prop-1-yn-1-yl)-3,4-dihydropyrimidin-1(2H)-yl)-3,4-dihydroxytetrahydrofuran-2-yl)methyl \\ ((2R,35,4R,5R)-5-(2,4-dioxo-5-(3-(6-(5-((3a5,45,6aR)-2-oxohexahydro-1H-thieno[3,4-d])midazol-4-yl)pentanamido) \\ ((2R,35,4R,5R)-5-(2,4-dioxo-5-(3-(6-(5-((3a5,45,6aR)-2-oxohexahydro-1H-thieno[3,4-d])midazol-4-yl)pentanamido) \\ ((3R,35,4R,5R)-5-(2,4-dioxo-5-(3-(6-(5-((3a5,45,6aR)-2-oxohexahydro-1H-thieno[3,4-d])midazol-4-yl)pentanamido) \\ ((3R,35,4R,5R)-5-(2,4-dioxo-5-(3-(6-(5-((3a5,45,6aR)-2-oxohexahydro-1H-thieno[3,4-d])midazol-4-yl)pentanamido) \\ ((3R,35,4R,5R)-5-(2,4-dioxo-5-(3-(6-(5-((3a5,45,6aR)-2-oxohexahydro-1H-thieno[3,4-d])midazol-4-yl)pentanamido) \\ ((3R,35,4R,5R)-5-(2,4-dioxo-5-(3-(6-(5-((3a5,45)-2-oxohexahydro-1H-thieno[3,4-d])midazol-4-yl)pentanamido) \\ ((3R,35,4R,5R)-5-(2,4-dioxo-5-(3-(6-(3a5,45)-2-oxohexahydro-1H-thieno[3,4-d])midazol-4-yl)pentanamido) \\ ((3R,35,4R)-3R)-(3R,35$

name: hydrogen triphospha

Solubility: good in water

Quality HPLC-MS (95%), testing in enzymatic reaction control:

Storage Storage: 12 months after receival at -20°C in the dark. Transportation: at room temperature for up to 3 weeks. Avoid excessive freeze-thaw cycles.