

---


## Biotin-X-hydrazide

<http://www.lumiprobe.com/p/biotin-aminohexanoic-acid-hydrazide>

Hydrazide biotin reagent with aminocaproic spacer useful for biotinylation of macromolecule reactive carbonyls.

Under acidic conditions, biotin-X-hydrazide couples selectively to aldehydes, and acid residues oxidized to aldehydes. It is used to prepare biotin-tagged glycoproteins, glycolipids, and antibodies via carboxylic groups cross-linking through carbodiimide chemistry. Hydrazide biotin is a reagent for common use in carbonylation research.

The aminocaproic spacer arm relieves steric hindrance for binding to streptavidin and avidins.



**Structure of Biotin-X-hydrazide**

---

### General properties

Appearance: white crystals

Molecular weight: 371.51

Molecular formula: C<sub>16</sub>H<sub>29</sub>N<sub>5</sub>O<sub>3</sub>S

Solubility: in DMSO, DMF

Quality control: NMR <sup>1</sup>H and HPLC-MS (95+%)

Storage conditions: 24 months after receipt at -20°C in the dark. Transportation: at room temperature for up to 3 weeks. Desiccate.

Legal statement: This Product is offered and sold for research purposes only. It has not been tested for safety and efficacy in food, drug, medical device, cosmetic, commercial or any other use. Supply does not express or imply authorization to use for any other purpose, including, without limitation, in vitro diagnostic purposes, in the manufacture of food or pharmaceutical products, in medical devices or in cosmetic products.