


Biotin-X-maleimide

<http://www.lumiprobe.com/p/biotin-aminohexanoic-acid-maleimide>

Biotin-X-maleimide is a thiol-reactive bifunctional linker for labeling SH groups. It can be used to attach biotin to proteins and peptides via cysteine residues and to other thiolated molecules, such as thiol-containing oligonucleotides.

Biotin-labeled compounds can then be linked to avidin or streptavidin for further purification or detection. The biotin group is relatively small and doesn't affect the biological activity of biotinylated molecules.

The reagent contains an aminocaproic spacer arm that relieves steric hindrance for binding the target molecule to streptavidin and avidin.



Structure of Biotin-X-maleimide

General properties

Appearance: beige powder

Molecular weight: 479.60

Molecular formula: $C_{22}H_{33}N_5O_5S$

Solubility: DMSO, DMF

Quality control: NMR 1H and HPLC-MS (95+%)

Storage conditions: 12 months after receipt at $-20^\circ C$ in the dark. Transportation: at room temperature for up to 3 weeks. Desiccate.

Legal statement: Product is offered and sold for research purposes only. Product is not tested for safety and efficacy in food, drug, medical device, cosmetic, no express or implied authorization to use for any other purpose, including, without limitation, in vitro diagnostic purposes, for humans or animals or for commercial purposes.